

REPORT DOCUMENTATION PAGE				Form Approved OMB NO. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA, 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>					
1. REPORT DATE (DD-MM-YYYY) 09-02-2013		2. REPORT TYPE Conference Proceeding		3. DATES COVERED (From - To) -	
4. TITLE AND SUBTITLE Using large-eddy simulation to investigate intermittency fluxes of clear-air radar reflectivity in the atmospheric boundary layer				5a. CONTRACT NUMBER W911NF-12-1-0089	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER 611102	
6. AUTHORS Andreas Muschinski, Peter P. Sullivan				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAMES AND ADDRESSES Northwest Research Associates, Inc. NorthWest Research Associates Inc. 4118 148th Ave. NE Redmond, WA 98052 -5164				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211				10. SPONSOR/MONITOR'S ACRONYM(S) ARO	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S) 61673-EV.6	
12. DISTRIBUTION AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other documentation.					
14. ABSTRACT Clear-air Doppler radars, also known as clear-air radar windprofilers, have been used for decades to remotely monitor wind velocities in the troposphere, stratosphere, and mesosphere. The traditional assumption is that the Doppler velocity (the first normalized moment of the Doppler spectrum) is an unbiased measure of the radial wind velocity within the radar's observation volume. Here we show that "intermittency fluxes", i.e., covariances of the turbulently fluctuating clear-air radar reflectivity and the turbulently fluctuating radial wind velocity, lead to					
15. SUBJECT TERMS Radar windprofiling, atmospheric turbulence, turbulence intermittency, Bragg scatter, clear-air radar, radio wave propagation					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	15. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Andreas Muschinski
a. REPORT UU	b. ABSTRACT UU	c. THIS PAGE UU			19b. TELEPHONE NUMBER 413-253-0129

Report Title

Using large-eddy simulation to investigate intermittency fluxes of clear-air radar reflectivity in the atmospheric boundary layer

ABSTRACT

Clear-air Doppler radars, also known as clear-air radar windprofilers, have been used for decades to remotely monitor wind velocities in the troposphere, stratosphere, and mesosphere. The traditional assumption is that the Doppler velocity (the first normalized moment of the Doppler spectrum) is an unbiased measure of the radial wind velocity within the radar's observation volume. Here we show that "intermittency fluxes", i.e., covariances of the turbulently fluctuating clear-air radar reflectivity and the turbulently fluctuating radial wind velocity, lead to systematic differences between the Doppler velocity and the (true) radial wind velocity. We use turbulent fields computationally generated by means of a large-eddy simulation to quantify this effect. We show that these biases may amount to several tens of centimeters per second in the atmospheric boundary layer, which is consistent with the biases observed with vertically pointing boundary-layer radar windprofilers.

Conference Name: 2013 IEEE International Symposium on Antennas and Propagation

Conference Date: July 07, 2013

Using large-eddy simulation to investigate intermittency fluxes of clear-air radar reflectivity in the atmospheric boundary layer

Andreas Muschinski¹ and Peter P. Sullivan²

¹ NorthWest Research Associates, Inc. (NWRA)
3380 Mitchell Lane, Boulder, CO 80301, andreas@cora.nwra.com

² Mesoscale and Microscale Meteorology Division,
National Center of Atmospheric Research, 3450 Mitchell Lane, Boulder, CO, 80307-3000

Submitted for presentation at
2013 IEEE International Symposium on Antennas and Propagation
(7-12 July 2013, Lake Buena Vista, Florida)

Using large-eddy simulation to investigate intermittency fluxes of clear-air radar reflectivity in the atmospheric boundary layer

Andreas Muschinski

NorthWest Research Associates, Inc.
Boulder, CO, United States
andreas@cora.nwra.com

Peter P. Sullivan

Mesoscale and Microscale Meteorology Division
National Center of Atmospheric Research
Boulder, CO, United States
pps@ucar.edu

Abstract—Clear-air Doppler radars, also known as clear-air radar windprofilers, have been used for decades to remotely monitor wind velocities in the troposphere, stratosphere, and mesosphere. The traditional assumption is that the Doppler velocity (the first normalized moment of the Doppler spectrum) is an unbiased measure of the radial wind velocity within the radar’s observation volume. Here we show that “intermittency fluxes”, i.e., covariances of the turbulently fluctuating clear-air radar reflectivity and the turbulently fluctuating radial wind velocity, lead to systematic differences between the Doppler velocity and the (true) radial wind velocity. We use turbulent fields computationally generated by means of a large-eddy simulation to quantify this effect. We show that these biases may amount to several tens of centimeters per second in the atmospheric boundary layer, which is consistent with the biases observed with vertically pointing boundary-layer radar windprofilers.

I. INTRODUCTION

Since the 1970s, clear-air radars, also known as radar windprofilers, have been used to remotely probe wind and turbulence characteristics in the troposphere, stratosphere, and mesosphere [1] [2] [3] [4] [5] [6]. Radar windprofilers exploit the very weak but still observable UHF and VHF radio waves backscattered from turbulent refractive-index perturbations in the optically clear atmosphere. The theory of acoustic and electromagnetic wave propagation through the turbulent atmosphere had been pioneered by Tatarskii [7] [8].

In the 1980s and 1990s, after radar windprofilers had reached a certain maturity, biases in the Doppler velocities observed with vertically pointing profilers were reported that could no longer be attributed to instrumental deficiencies. Instead, evidence accumulated that such biases were caused by certain correlations between wind velocity fluctuations and reflectivity fluctuations [9] [10]. Those mechanisms, however, relied on the presence of gravity waves [9] or Kelvin-Helmholtz billows [10] and therefore could not account for the velocity biases of a few tens of centimeters per second that Angevine [11] observed within the convective boundary layer. Observational and theoretical aspects of those Doppler velocity biases have been discussed in more detail in [12] and [6].

Around 1970, the large-eddy simulation (LES) technique was pioneered by Lilly [13] and Deardorff [14]. LES is now

the method of choice for computationally investigating the structure and dynamics of the atmospheric boundary layer, and LES experiments may be considered in many respects to be equivalent, if not superior, to real-world boundary-layer experiments [15].

In 1999, we used for the first time LES-generated fields to synthesize realistic clear-air radar signals [16]. The main motivation for that study was to reproduce Angevine’s [11] observations of velocity biases in the convective boundary layer.

II. INTERMITTENCY IN ATMOSPHERIC TURBULENCE

Classical turbulence theory [17] [18] and the classical theory of wave propagation through the turbulent atmosphere [7] [8] rely on the assumption that turbulent dissipation rates and turbulent structure parameters may be treated as deterministic quantities that vary only slowly in space and time. In their two 1962 landmark papers, however, Kolmogorov [19] and Obukhov [20] showed that at high Reynolds numbers as encountered in the atmosphere and the oceans, dissipation rates have to be treated as random variables and that the statistics of that intermittency has to be accounted for in a complete theory of locally homogeneous turbulence.

III. INTERMITTENCY FLUXES AND DOPPLER VELOCITY BIASES

The Doppler velocity v_D is the first moment M_1 of the Doppler spectrum, normalized by the zeroth moment. Assuming that the profiler points in the vertical direction and interpreting the Doppler spectrum as a histogram of reflectivity-weighted radial (here vertical) wind velocities, we can write

$$v_D = \frac{M_1}{M_0} = \frac{\langle \eta w \rangle}{\langle \eta \rangle}, \quad (1)$$

where w is the vertical wind velocity and η is the clear-air volume reflectivity. Now, we treat η and w as random variables and write them as sums of mean value and fluctuation:

$$\eta = \langle \eta \rangle + \eta', \quad w = \langle w \rangle + w'. \quad (2)$$

This gives

$$v_D = \frac{\langle \eta \rangle \langle w \rangle + \langle \eta' w' \rangle}{\langle \eta \rangle} = \langle w \rangle + \Delta w, \quad (3)$$

where

$$\Delta w = \frac{\langle \eta' w' \rangle}{\langle \eta \rangle} \quad (4)$$

is the velocity bias.

Note that $\langle \eta' w' \rangle$ may be interpreted as a turbulent clear-air reflectivity flux. We suggest to refer to such a flux as *intermittency flux* because in traditional propagation theory [7] [8], which does not account for intermittency and which treats η (which is proportional to C_n^2 [7]) as a deterministic variable, $\langle \eta' w' \rangle$ is always zero.

IV. USING LARGE-EDDY SIMULATION TO COMPUTE INTERMITTENCY FLUXES

We used a large-eddy simulation similar to the one in our previous study [16], where we assumed that the dwell time is short compared to the large-eddy overturning time scale and that the radar's observation volume is small compared to the energy-containing scales in the convective boundary layer. In the previous study, we did not find a significant velocity bias. In the current study, however, we allowed the effective radar sampling volume to be large compared to the large-eddy scale, and we find velocity biases with magnitudes of up to 50 cm s^{-1} (see Fig. 1), consistent with Angevine's [11] observations.




Fig. 1. Vertical profiles of Doppler velocity biases caused by intermittency fluxes of the temperature structure parameter, C_T^2 , and the humidity structure parameter, C_q^2 , respectively. The profiles were computed from an LES-generated convective boundary layer similar to the LES described in our previous study [16].

V. SUMMARY AND CONCLUSIONS

“Intermittency fluxes” of clear-air radar reflectivity can lead to Doppler velocity biases observed with vertically pointing radar windprofilers in the convective boundary layer (CBL).

We have used the large-eddy simulation (LES) technique to computationally generate a CBL, and we computed vertical profiles of Doppler velocity biases that agree qualitatively and quantitatively with biases observed in field measurements.

REFERENCES

- [1] R. F. Woodman and A. Guillén, “Radar observations of winds and turbulence in the stratosphere and mesosphere,” *J. Atmos. Sci.*, vol. 31, pp. 493–505, 1974.
- [2] K. S. Gage, “Radar observations of the free atmosphere: structure and dynamics,” in *Radar in meteorology*, D. Atlas, Ed. Boston: Amer. Meteor. Soc., 1990, pp. 534–565.
- [3] J. Röttger and M. F. Larsen, “UHF/VHF radar techniques for atmospheric research and wind profiler applications,” in *Radar in meteorology*, D. Atlas, Ed. Boston: Amer. Meteor. Soc., 1990, pp. 235–281.
- [4] E. E. Gossard, “Radar research on the atmospheric boundary layer,” in *Radar in meteorology*, D. Atlas, Ed. Boston: Amer. Meteor. Soc., 1990, pp. 477–527.
- [5] J. M. Wilczak, E. E. Gossard, W. D. Neff, and W. L. Eberhard, “Ground-based remote sensing of the atmospheric boundary layer: 25 years of progress,” *Boundary-Layer Meteorol.*, vol. 78, pp. 321–349, 1996.
- [6] A. Muschinski, V. Lehmann, L. Justen, and G. Teschke, “Advanced radar wind profiling,” *Meteorol. Z.*, vol. 14, pp. 609–625, 2005.
- [7] V. I. Tatarskii, *Wave propagation in a turbulent medium*. New York: McGraw-Hill, 1961.
- [8] —, *The effects of the turbulent atmosphere on wave propagation*. Jerusalem, Israel: Israel Program for Scientific Translation, 1971.
- [9] G. D. Nastrom and T. E. VanZandt, “Mean vertical motion seen by radar wind profilers,” *J. Appl. Meteor.*, vol. 33, pp. 984–995, 1994.
- [10] A. Muschinski, “Possible effect of Kelvin-Helmholtz instability on VHF radar observations of the mean vertical wind,” *J. Appl. Meteor.*, vol. 35, pp. 2210–2217, 1996.
- [11] W. M. Angevine, “Errors in mean vertical velocities measured by boundary layer wind profilers,” *J. Atmos. Oceanic Technol.*, vol. 14, pp. 565–569, 1997.
- [12] A. Muschinski, “Local and global sampling of clear-air Doppler radar signals,” *Radio Sci.*, vol. 39, pp. RS1008, doi:10.1029/2003RS002908, 2004.
- [13] D. K. Lilly, “The representation of small-scale turbulence in numerical simulation experiments,” in *Proc. IBM Sci. Comput. Symp. Environm. Sci. (Yorktown Heights, Nov. 14th to 15th 1966)*. Yorktown Heights, N. Y.: Thomas J. Watson Res. Center, 1967, pp. 195–210.
- [14] J. W. Deardorff, “Three-dimensional numerical modeling of the planetary boundary layer,” in *Workshop on micrometeorology*, D. A. Haugen, Ed. Boston: Amer. Meteor. Soc., 1973, pp. 271–311.
- [15] J. C. Wyngaard, *Turbulence in the atmosphere*. Cambridge, England: Cambridge University Press, 2010.
- [16] A. Muschinski, P. P. Sullivan, D. B. Wuertz, R. J. Hill, S. A. Cohn, D. H. Lenschow, and R. J. Doviak, “First synthesis of wind-profiler signals on the basis of large-eddy simulation data,” *Radio Sci.*, vol. 34, pp. 1437–1459, 1999.
- [17] A. N. Kolmogorov, “Local structure of turbulence in an incompressible fluid at very high Reynolds numbers,” *Dokl. Akad. Nauk SSSR*, vol. 30, pp. 299–303, 1941.
- [18] G. K. Batchelor, *The theory of homogeneous turbulence*. Cambridge: Cambridge University Press, 1953.
- [19] A. N. Kolmogorov, “A refinement of previous hypotheses concerning the local structure of turbulence in a viscous incompressible fluid at high Reynolds number,” *J. Fluid Mech.*, vol. 13, pp. 82–85, 1962.
- [20] A. M. Obukhov, “Some specific features of atmospheric turbulence,” *J. Fluid Mech.*, vol. 13, pp. 77–81, 1962.